

**Proposed Renewstable<sup>®</sup> Sivutse and Associated  
Infrastructure near the Majuba Power Station,  
Mpumalanga Province**

**Preliminary Avifaunal Assessment &  
Site Sensitivity Verification**



Compiled for



By



**Cossypha**  
Ecological

July 2024

## REPORT PRODUCTION

Specialist	Role	Project Component	Qualifications and Professional Registration
Robyn Phillips	Terrestrial Ecologist (Avifaunal Specialist)	Field work (data collection) and report compilation	MSc (Zoology) UNP SACNASP: <i>Pr.Sci.Nat.</i> 400401/12 Fields: Zoological and Ecological

## CONTACT INFORMATION

Robyn Phillips  
16 MacDonald Road  
Woodside, Westville  
KwaZulu-Natal, 3629  
Cell: 084 695 1648  
Email: robyn@cosypha.co.za

## SPECIALIST DECLARATION OF INDEPENDENCE

I, **Robyn Phillips**, in my capacity as a specialist consultant, hereby declare that I –

- Act as an independent consultant;
- Do not have any financial interest in the undertaking of the activity, other than remuneration for the work performed in terms of the National Environmental Management Act, 1998 (Act No. 107 of 1998);
- Do not have and will not have vested interest in the proposed activity proceeding;
- Have no, and will not engage in, conflicting interests in the undertaking of the activity;
- Undertake to disclose, to the Competent Authority, any material information that has or may have the potential to influence the decision of the competent authority or the objectivity of any report, plan or document required in terms of the National Environmental Management Act, 1998 (Act 107 of 1998);
- Will provide the Competent Authority with access to all information at my disposal regarding the application, whether such information is favourable to the applicant or not;
- As a registered member of the South African Council for Natural Scientific Professions, will undertake my profession in accordance with the Code of Conduct of the Council, as well as any other societies to which I am a member;
- Based on information provided to me by the project proponent and in addition to information obtained during the course of this study, have presented the results and conclusion within the associated document to the best of my professional ability;
- Reserve the right to modify aspects pertaining to the present investigation should additional information become available through ongoing research and/or further work in this field; and
- Undertake to have my work peer reviewed on a regular basis by a competent specialist in the field of study for which I am registered.

July 2024

---

**Robyn Phillips** *Pr.Sci.Nat.*  
Terrestrial Ecologist  
SACNASP Reg. No. 400401/12

---

Date

## TABLE OF CONTENTS

Report Production .....	ii
Contact Information .....	ii
Specialist Declaration of Independence.....	ii
Table of Contents.....	iii
List of Figures.....	iv
List of Tables.....	iv
Abbreviations .....	v
<b>1. Introduction and Project Description .....</b>	<b>1</b>
1.1. Project Description .....	1
1.2. Location.....	2
1.3. Reporting Requirements .....	2
1.4. Terms of Reference.....	4
<b>2. Methodology .....</b>	<b>4</b>
2.1. Stage 1: Preliminary Assessment.....	5
2.2. Assumptions and Limitations .....	6
<b>3. Desktop Assessment Results .....</b>	<b>6</b>
3.1. Description of the Study Area .....	6
3.2. Distribution of Avifauna in the Study Area.....	10
<b>4. Field Results.....</b>	<b>14</b>
4.1. Site Description.....	14
4.2. Avifaunal Habitats in the Study Area and Surrounds .....	17
4.3. Bird Species Occurrence in the Study Area.....	18
4.3.1. Bird Observations .....	18
4.3.2. Birds of Conservation Concern .....	19
4.3.3. Priority Species .....	20
4.4. Key Habitats and Preliminary Site Sensitivity.....	21
<b>5. Site Sensitivity Verification.....</b>	<b>24</b>
<b>6. Preliminary Impacts .....</b>	<b>24</b>
<b>7. Pre-construction Monitoring Requirements.....</b>	<b>25</b>
<b>8. Recommendations .....</b>	<b>27</b>
<b>9. References.....</b>	<b>27</b>
<b>10. Appendices .....</b>	<b>29</b>
APPENDIX A: Abridged CV of the Specialist .....	29

## LIST OF FIGURES

Figure 1: Example of a hydrogen power plant with solar PV arrays .....	1
Figure 2: Location of the Sivutse study area .....	2
Figure 3: DFFE Screening Tool map of relative sensitivity for the Animal Species theme for the Sivutse A site	3
Figure 4: DFFE Screening Tool map of relative sensitivity for the Animal Species theme for the Sivutse B site	3
Figure 5: Historical aerial images of Sivutse A (left) and Sivutse B (right) from 1957 .....	7
Figure 6: Aerial overview of the Sivutse study area and surrounds .....	8
Figure 7: The Sivutse study area in relation to national Protected Areas .....	9
Figure 8: Habitat features of the Sivutse study area .....	16
Figure 9: Preliminary avifaunal habitat sensitivity of the Sivutse study area .....	23

## LIST OF TABLES

Table 1: Avifaunal priority species occurring within pentads 2700_2945, 2700_2950, 2705_2945, and 2705_2950 including Reporting Rate (RR) .....	10
Table 2: Species recorded in the study area and surroundings listed in taxonomic order. SCC are highlighted in red.....	20
Table 3: Summary of sensitivity categories.....	22
Table 4: Possible impacts arising from the proposed development.....	25
Table 5: Recommended avifaunal assessment regimes (Jenkins <i>et al.</i> , 2017) .....	26

## ABBREVIATIONS

BA	Basic Assessment
BESS	Battery Energy Storage System
BIRP	Birds in Reserves Project
CAR	Co-ordinated Avifaunal Road Counts
CWAC	Co-ordinated Wetland Counts
DEA	Department of Environmental Affairs (now DFFE)
DFFE	Department of Forestry, Fisheries and the Environment
EA	Environmental Authorisation
EAP	Environmental Assessment Practitioner
EIA	Environmental Impact Assessment
EN	Endangered
EWT	Endangered Wildlife Trust
GBIF	Global Biodiversity Information Facility
GN	General Notice
IUCN	International Union for Conservation of Nature
kV	Kilovolts
LC	Least Concern
MW	Megawatt
NEMA	National Environmental Management Act 107 of 1998
NT	Near Threatened
PV	Photovoltaic
QDGC	Quarter Degree Grid Cell
SABAP	South African Bird Atlas Project
SANBI	South African National Biodiversity Institute
SCC	Species of Conservation Concern
SEF	Solar Energy Facility
SPC	Special Purpose Company
VU	Vulnerable

## 1. INTRODUCTION AND PROJECT DESCRIPTION

Hydrogene de France (HDF), under its Special Purpose Company (SPC) Renewstable<sup>®</sup> Mpumalanga (Pty) Ltd, proposes to develop the Renewstable<sup>®</sup> Sivutse renewable power plant near the Majuba Power Station in Mpumalanga. The project is part of a cluster of similar developments, which are high-capacity renewable power plants based on hydrogen energy storage technology. Nsovo Environmental Consulting (Pty) Ltd was appointed to undertake an Environmental Impact Assessment (EIA) process required in terms of the National Environmental Management Act 107 of 1998 (NEMA), and the Environmental Impact Assessment (EIA) Regulations, 2014 (GN R982, as amended by GN R326) for the project. Cossypha Ecological was appointed to conduct an Avifaunal Impact Assessment for the proposed development to inform the EIA process.

### 1.1. PROJECT DESCRIPTION

A development area of approximately 437 ha, split between two parcels of land known as Bergvliet (hereafter referred to as Sivutse A) and Riet C (hereafter referred to as Sivutse B), has been earmarked for the proposed Renewstable<sup>®</sup> Sivutse project. This facility will provide between 12 MW and 55 MW of electricity services to the country daily over a period of at least 25 years from the commissioning of the plant. The cluster of plants are scheduled to be commissioned in 2027 and will contribute to the greening of the local power grid and enhance the territory's energy independence.

The Renewstable<sup>®</sup> power plants convert the electricity from a photovoltaic (PV) solar park into hydrogen through an electrolyser system, then stores this hydrogen in compressed gas form, and restitutes the electricity to the grid through a fuel cell system when the PV park no longer produces enough energy. Hydrogen technologies rely on the electrochemical properties of water by decomposing and then recomposing a water molecule (H<sub>2</sub>O) using electrical energy, without emitting greenhouse gases. Therefore, the system does not generate any harmful atmospheric emissions, only oxygen, with traces of water as vapor, hydrogen, and nitrogen during the maintenance phase. The site would also include battery power storage to maximise plant performance and improve customer service. A Battery Energy Storage System (BESS) provides the end-of-the-day peak power and, in combination with the hydrogen storage, ensures the stability of the electricity service. In the Renewstable<sup>®</sup> Power Plant BESS is complementary to hydrogen being used as short-term energy storage and for power regulation. There are many technologies on the BESS market, but the current leading technology and more suitable for Renewstable<sup>®</sup> currently is the Lithium-Ion (Li-Ion) BESS.




Figure 1: Example of a hydrogen power plant with solar PV arrays

## 1.2. LOCATION

The Renewable<sup>®</sup> Sivutse project area is split between two land parcels, the Sivutse A site located ~7 km to the east of the Majuba Power Station and the Sivutse B site located approximately 4 km to the north-east of the Majuba Power Station. Together these will be referred to as the study area. The study area is situated ~8 km south of the town of Amersfoort within the Dr Pixley Ka Isaka Seme Local Municipality in the Gert Sibande District of Mpumalanga Province (**Figure 2**). Overall, the study area is 437 ha in extent with the Sivutse A site (130 ha) occurring on a portion of the Farm Bergvliet 65-HS Ptn 34, and the Sivutse B site (307 ha) occurring on a portion of the Farm Rietfontein 66-HS Ptn 4. The study area falls within Quarter Degree Grid Cell (QDGC) 2729BB and lies between 27°04'23.7" and 27°06'02.2" south and 29°47'51.16" and 29°51'34.6" east. The study area is gently undulating with a range in altitude from around 1 676 to 1 741 m above mean sea level (a.m.s.l.).




Figure 2: Location of the Sivutse study area

## 1.3. REPORTING REQUIREMENTS

Environmental site sensitivity reports were generated for each site for the project on the 26<sup>th</sup> of June 2024 using the Department of Forestry, Fisheries, and the Environment (DFFE) National Web-Based Environmental Screening Tool. For the proposed development sites, the Screening Tool Report identified a possible site environmental sensitivities of **High** and **Medium** for Aves under the Animal Species theme due to the potential occurrence of the the following bird Species of Conservation Concern (SCC) within the study area (**Figure 4** and **Figure 4**):

- High: *Circus ranivorus* African Marsh Harrier (Endangered (EN))
- High: *Spizocorys fringillaris* Botha's Lark (EN)
- High: *Sagittarius serpentarius* Secretarybird (Vulnerable (VU))
- High: *Geronticus calvus* Southern Bald Ibis (VU)
- Medium: *Eupodotis senegalensis* White-bellied Bustard (VU)
- Medium: *Sagittarius serpentarius* Secretarybird (VU)

- Medium: *Tyto Capensis* African Grass Owl (VU)
- Medium: *Circus ranivorus* African Marsh Harrier (EN)
- Medium: *Spizocorys fringillaris* Botha’s Lark (EN)
- Medium: *Neotis denhami* Denham’s Bustard (VU)
- Medium: *Balearica regulorum* Grey Crowned Crane (EN)
- Medium: *Geronticus calvus* Southern Bald Ibis (VU)

MAP OF RELATIVE ANIMAL SPECIES THEME SENSITIVITY

MAP OF RELATIVE AVIAN THEME SENSITIVITY




Figure 3: DFFE Screening Tool map of relative sensitivity for the Animal Species theme (left) and Avian theme (right) for the Sivutse A site

MAP OF RELATIVE ANIMAL SPECIES THEME SENSITIVITY

MAP OF RELATIVE AVIAN THEME SENSITIVITY




Figure 4: DFFE Screening Tool map of relative sensitivity for the Animal Species theme (left) and Avian theme (right) for the Sivutse B site

Therefore, based on the environmental sensitivities of the proposed development footprint, the screening tool suggested the following applicable specialist assessment for inclusion in the EIA report:

- Avian Impact Assessment

The assessment must be compiled in accordance with the requirements of the *Procedures for the Assessment and Minimum Criteria for Reporting on Identified Environmental Themes when Applying for EA* (GN R320 of 2020) and comply with the following gazetted protocol, which replaces the requirements of Appendix 6 of the EIA Regulations, 2014 (as amended) in terms of NEMA:

- *Protocol for the Specialist Assessment and Minimum Report Content Requirements for Environmental Impacts on Animal Species* (GN 1150 of 30 October 2020) as amended 28 July 2023.

According to the above-mentioned protocol, the report must follow the Species Environmental Assessment Guidelines (SANBI, 2020), which prescribes the Best Practice Guidelines: Birds & Solar Energy (Jenkins *et al.*, 2017) for assessing and monitoring the impact of solar power generating facilities on birds in southern Africa, established by BirdLife South Africa and the Wildlife and Energy Programme of the Endangered Wildlife Trust (EWT).

According to the above mentioned gazetted protocols, prior to commencing with a specialist assessment, the current use of the land and the potential environmental sensitivity of the site identified by the screening tool must be confirmed by undertaking a **Site Sensitivity Verification (SSV)**. The purpose of the SSV is to confirm the actual use of the land on the ground versus that which has been identified by the screening tool. The SSV must confirm or refute the need to employ the various specialists as identified in the screening report. The SSV conducted for this study confirmed both the Animal Species theme and the Avian theme to be of **Medium** sensitivity for the Sivutse A site, and **High** sensitivity for the Sivutse B site due to the presence of bird SCC as well as a large area of connected natural grassland habitat for these species in the study area. **Refer to further information provided in the sections that follow and the SSV in Section 5.**

**This Report covers Stage 1 of the methodology prescribed by the Best Practice Guidelines (Jenkins *et al.*, 2017) and entails the Preliminary Avifaunal Assessment as well as the SSV of the DFFE Environmental Screening Report outcomes.**

#### 1.4. TERMS OF REFERENCE

The Terms of Reference for this report were to:

- Undertake a preliminary field survey of the study area to identify and map areas of opportunity and constraint within the property to inform the layout.
- Compile a photographic record of the characteristics of the study area, including major habitats and sensitive areas.
- Provide a verification of the site sensitivities identified by the DFFE screening tool (SSV Report).
- Compile a Preliminary Avifaunal Assessment Report that provides an overview of the ecological context, likely impacts, and potential red flags to development, from an avifaunal perspective, covering **Stage 1** of the methodology prescribed by the Guidelines (Jenkins *et al.*, 2017); and
- Provide maps and shapefiles based on the preliminary findings.

## 2. METHODOLOGY


The solar energy industry is expanding rapidly in southern Africa, and the nature and implications of potential negative effects on birds, through the destruction of habitat, the displacement of populations from preferred habitat, and collision and burn mortality associated with the solar infrastructure, are poorly understood. To fully understand and avoid and minimise the possible impacts of solar energy on the region's birds, it is essential that sufficient, project- and site-specific data are gathered to both inform the avifaunal impact assessment process and build our understanding of the impacts and potential mitigation measures (Jenkins *et al.*, 2017). According to Jenkins *et al.* (2017), an avifaunal impact assessment for Solar Energy Facilities (SEFs) must follow a tiered process that follows pre-determined stages depending on the conditions of the site:

**Stage 1 – Preliminary Assessment:** part of planning for an EIA application (i.e. pre-application). This provides an overview of the ecological context, likely impacts and potential red flags to development, identify alternatives and determine the appropriate assessment regime.

**Stage 2 – Data Collection:** an in-depth study including structured and repeated data collection on which to base the impact assessment report and provide a baseline against which post-construction monitoring can be compared.

**Stage 3 – Impact Assessment:** informed by the data collected during Stage 2.

**Stage 4 – Monitoring and Mitigation:** during construction and post-construction monitoring to inform mitigation, informed by the data collected during Stage 2 (regime 2 and 3 only).



*This document reports information and results for stage 1.*

## 2.1. STAGE 1: PRELIMINARY ASSESSMENT

According to Jenkins *et al.* (2017) the preliminary assessment should yield a preliminary avifaunal assessment report, which describes the relative sensitivity of the study area, highlights any red flags to development, and determines whether additional baseline data collection is necessary to fully inform the Avifaunal Impact Assessment Report. The preliminary assessment is based on desk-top review and a site survey conducted over the study area in the summer season from the from the 14<sup>th</sup> to the 16<sup>th</sup> of November 2023. The findings are incorporated into a report aimed to characterise the study area in terms of habitats present, the overall site sensitivity, and delineate areas that are potentially highly sensitive and no-go areas that may need to be avoided by the development. The sensitivity analysis follows the methodology prescribed in the Species Environmental Assessment Guideline (SANBI, 2020). Preliminary assessment of impacts and general recommendations are also provided.

Prior to the site visit, a comprehensive list of bird species occurring in the area was compiled using electronic databases within Roberts VII Multimedia Birds of Southern Africa (SA Birding, 2011) where distribution maps have been interpreted and updated from the Atlas of Southern African Birds (Harrison *et al.*, 1997). The search was confined to the quarter degree grid cell (QDGC) in which the study area falls (i.e. atlas area of 15' × 15' – roughly 24 × 27 km) to get a comprehensive list of species for the region. The data was supplemented with current Southern African Bird Atlas Project 2 (SABAP2, 2022) data, which is recorded per pentad (a 5' × 5' coordinate spatial grid reference – one QDGC comprises of nine pentads). Species of conservation concern (SCC) that could potentially occur in the greater study area were noted and their habitat requirements determined by consulting the relevant literature. Bird names follow the International Ornithological Congress (IOC) World Bird

List (v13.2) (Gill *et al.*, 2023) while conservation status follows the latest Red Data Book of Birds (Taylor *et al.*, 2015). Other online databases such as Co-ordinated Wetland Counts (CWAC), Co-ordinated Avifaunal Road Counts (CAR), Birds in Reserves Project (BIRP), Global Biodiversity Information Facility (GBIF), and iNaturalist were searched for avifaunal SCC potentially occurring in the area.

Prior to the site visit, recent and historical aerial imagery using Google Earth and the Chief Directorate National Geospatial Information (CDNGI) Geospatial Portal (<http://www.cdngiportal.co.za/cdngiportal/>) was reviewed to differentiate areas with natural vegetation versus modified and transformed areas of the study area. Available online databases relating to regional biodiversity conservation planning, e.g. national vegetation types, threatened ecosystems, the relevant provincial spatial conservation or biodiversity plan, Important Bird Areas (IBAs), and National Protected Areas etc. were also reviewed with the aim of flagging any potentially important areas of the site that would need special attention during the site visit.

Survey techniques included on-site meander searches, observations for priority species, and focussed counts at habitats such as wetlands, dams, and koppies. During meander searches through the study area, changes in land cover and habitat, as well as avifauna present in the study area were observed and recorded. Landscape features that were considered of high ecological importance were mapped.

## 2.2. ASSUMPTIONS AND LIMITATIONS

The following assumptions and limitations pertain to this report:

- Habitat boundaries usually consist of subtle transitional zones or ecotones, which cannot be captured as distinct lines. Boundaries of habitat types are therefore approximately defined.
- Habitat types are defined and mapped in the context of use by birds and not in terms of botanical species associations. Similarly, the riparian habitat associated with rivers, wetlands, and dams are defined in terms of broad habitat use by birds and do not denote the boundaries of wetlands and watercourses.
- The preliminary assessment was conducted over the entire ~437 ha study area comprised of the two sites to get an overview of habitats, landscape features, and sensitivities. While it's unlikely, any of the habitat delineations are subject to change if new sensitivities come to light following the more detailed seasonal assessments.

## 3. DESKTOP ASSESSMENT RESULTS

### 3.1. DESCRIPTION OF THE STUDY AREA

The study area lies in the eastern highveld parts of the country, within the Grassland Biome and the Mesic Highveld Grassland Bioregion (Rutherford and Westfall, 1994; Mucina and Rutherford, 2006), which is characterised by cold dry winters and mild summers. Rain falls mostly in summer with a Mean Annual Precipitation (MAP) of 694 mm. The highest rainfall occurs in January and the lowest falls in June / July. Maximum temperatures reach around 27°C in summer and minimum temperatures drop to around 8°C in winter. Incidence of frost is very high (Mucina and Rutherford, 2006).

According to SANBI (2018) the study area is situated within the Amersfoort Highveld Clay Grassland vegetation type, which is endemic to South Africa and occurs in KwaZulu-Natal and Mpumalanga Provinces. Amersfoort Highveld Clay Grassland consists of undulating grassland plains, with scattered patches of dolerite outcrops. The grassland is characteristically short and closed, dominated by *Themeda triandra* and is often severely grazed (Mucina and Rutherford, 2006; SANBI, 2021). Amersfoort Highveld Clay Grassland is classified as Least Concern,

with a target of 27%, only 3.6% is statutorily conserved, with around 45% transformed mainly for agriculture (Mucina and Rutherford, 2006; SANBI, 2021; DFFE, 2022).

The landscape of the Sivutse study area is rural in nature occurring in farmland used mostly for cattle grazing and cultivation. The Sivutse A site is situated within a fenced Eskom property (the old Majuba Colliery underground mine) and comprises mostly disturbed grassland used for grazing cattle, scattered with clumps of alien trees. Derelict buildings (the old mine administrative buildings) and a small dam occur towards the centre of the site. The site once comprised natural grassland (**Figure 5**) but was converted over 40 years ago. The Sivutse B site is comprised of natural open grassland vegetation interspersed with natural drainage lines and small farm dams, as well as cultivated and fallow fields. The natural grassland vegetation in the northern section of the site forms part of a relatively continuous expanse of open grassland found in the surrounding areas that is relatively undisturbed and according to historical aerial imagery (from 1957 and 1964 (**Figure 5**) to today), has not been modified by ploughing in the last ~67 years and has likely only been used for grazing. The surrounding areas comprise privately owned farmland with natural open grassland, scattered with cultivated fields, natural drainage lines, and small farm dams. The Majuba Power Station occurs ~2.5 km to the west of Sivutse B and the N11 highway occurs ~2 km to the east of Sivutse A (**Figure 6**).



**Figure 5: Historical aerial images of Sivutse A (left) from 1964 and Sivutse B (right) from 1957**

According to the Mpumalanga Biodiversity Sector Plan (MTPA, 2014), the northern section of the Sivutse B site and the southern section of the Sivutse A site is comprised of Critical Biodiversity Area (CBA) Irreplaceable mostly associated with the natural grassland vegetation. The far western section is classified as “Other Natural Area” with a strip of Ecological Support Area (ESA). The remainder, mostly cultivated fields, is classified as “Heavily Modified”. The Majuba Nature Reserve occurs ~700 m to the west of the Sivutse B site and the Afrikan Farms Protected Environment occurs ~18.6 km to the east of the Sivutse A site. The study area falls between two Important Bird Areas (IBAs), the Grasslands IBA (ZA016) occurs ~2.2 km to the south and ~6 km to the east, while the Amersfoort-Bethal-Carolina IBA (ZA014) occurs ~7.4 km to the north of the site. In addition, most of the remaining natural grassland on the site and in the surrounding areas is flagged as Protected Area Expansion Priority Areas (**Figure 7**).




Figure 6: Aerial overview of the Sivutse study area and surrounds




Figure 7: The Sivutse study area in relation to national Protected Areas

### 3.2. DISTRIBUTION OF AVIFAUNA IN THE STUDY AREA

The region is relatively high in avifaunal diversity with around 308 bird species known to occur within the QDGC (an atlas area of 15' × 15' – roughly 24 × 27 km) that the study area falls within (2729BB), according to the distribution maps in Roberts VII Multimedia Birds of Southern Africa (SA Birding, 2011). Approximately 87% of the total species in the QDGCs are associated with grassland habitat, farmlands, and inland water habitats, which is the character of the study area. This demonstrates that the available habitats within the study area are able to support the majority of bird species found within the QDGC.

The Southern African Bird Atlas Project (SABAP2) has been collecting data since 2007 and includes data from the previous SABAP1 (1987-1991). SABAP2 aims to map the distribution and relative abundance of birds in southern Africa. SABAP2 data is recorded per pentad (a 5' × 5' coordinate spatial grid reference and a subset of the QDGC – one QDGC comprises of nine pentads. 5' × 5' = roughly 8 × 9 km) and therefore represents a more focussed search. Reporting rates are expressed as a percentage of the number of times a species was seen in a pentad divided by the number of times the pentad was surveyed. According to SABAP2 data, 156 bird species have been recorded in the pentads in which the study area falls (pentads 2700\_2945, 2700\_2950, 2705\_2945, and 2705\_2950). This includes 13 species of conservation concern (SCC) (see species highlighted in **Table 1**).

Priority species in terms of sensitivity to solar PV development impacts include any Red List (SCC), range-restricted species, species that congregate in large numbers (gregarious species), and large-bodied species such as waterfowl, herons, gamebirds, and raptors (including owls and vultures) (Jenkins *et al.*, 2017). **Table 1** lists priority species that have been recorded within the pentads, with the SABAP2 reporting rate. The higher the reporting rate, the higher the likelihood of the species occurring in the study area if suitable habitat exists. A reporting rate of zero implies that the bird was recorded with an ad-hoc sighting. See **Table 2** in **Section 4.3.3** for species recorded in the study area.

**Table 1: Avifaunal priority species occurring within pentads 2700\_2945, 2700\_2950, 2705\_2945, and 2705\_2950 including Reporting Rate (RR)**

Common Name	Scientific Name	Priority Species	Threat Status (RSA / IUCN)	SABAP2 RR (%)
Blacksmith Lapwing	<i>Vanellus armatus</i>		LC / LC	100
Hadada Ibis	<i>Bostrychia hagedash</i>	Gamebird	LC / LC	100
African Stonechat	<i>Saxicola torquatus</i>		LC / LC	100
Cape Longclaw	<i>Macronyx capensis</i>		LC / LC	100
Southern Masked Weaver	<i>Ploceus velatus</i>	Gregarious	LC / LC	100
Southern Red Bishop	<i>Euplectes orix</i>	Gregarious	LC / LC	100
Helmeted Guineafowl	<i>Numida meleagris</i>	Gamebird	LC / LC	87.5
Ring-necked Dove	<i>Streptopelia capicola</i>		LC / LC	87.5
Black-winged Kite	<i>Elanus caeruleus</i>	Raptor	LC / LC	87.5
Black-headed Heron	<i>Ardea melanocephala</i>	Gamebird	LC / LC	87.5
Western Cattle Egret	<i>Bubulcus ibis</i>	Waterfowl	LC / LC	87.5
Levaillant's Cisticola	<i>Cisticola tinniens</i>		LC / LC	87.5
Red-capped Lark	<i>Calandrella cinerea</i>		LC / LC	87.5
Southern Grey-headed Sparrow	<i>Passer diffusus</i>	Gregarious	LC / LC	87.5
African Pipit	<i>Anthus cinnamomeus</i>		LC / LC	87.5
Long-tailed Widowbird	<i>Euplectes progne</i>	Gregarious	LC / LC	87.5
Egyptian Goose	<i>Alopochen aegyptiaca</i>	Waterfowl	LC / LC	75
Reed Cormorant	<i>Microcarbo africanus</i>	Waterfowl	LC / LC	75

Common Name	Scientific Name	Priority Species	Threat Status (RSA / IUCN)	SABAP2 RR (%)
Barn Swallow	<i>Hirundo rustica</i>		LC / LC	75
Cape Sparrow	<i>Passer melanurus</i>	Gregarious	LC / LC	75
Cape Wagtail	<i>Motacilla capensis</i>		LC / LC	75
Quailfinch	<i>Ortygospiza atricollis</i>	Gregarious	LC / LC	75
Rock Dove	<i>Columba livia</i>		LC / LC	63.6
Swainson's Spurfowl	<i>Pternistis swainsonii</i>	Gamebird	LC / LC	62.5
Yellow-billed Duck	<i>Anas undulata</i>	Waterfowl	LC / LC	62.5
Blue Korhaan	<i>Eupodotis caerulescens</i>	SCC	LC; SLS / NT	62.5
Red-knobbed Coot	<i>Fulica cristata</i>	Waterfowl	LC / LC	62.5
Amur Falcon	<i>Falco amurensis</i>	Raptor	LC / LC	62.5
African Sacred Ibis	<i>Threskiornis aethiopicus</i>	Waterfowl	LC / LC	62.5
Southern Fiscal	<i>Lanius collaris</i>		LC / LC	62.5
Ant-eating Chat	<i>Myrmecocichla formicivora</i>		LC / LC	62.5
Banded Martin	<i>Riparia cincta</i>		LC / LC	62.5
South African Cliff Swallow	<i>Petrochelidon spilodera</i>	Gregarious	LC; NE / LC	62.5
Common Waxbill	<i>Estrilda astrild</i>	Gregarious	LC / LC	62.5
Cape Canary	<i>Serinus canicollis</i>	Gregarious	LC / LC	62.5
Spur-winged Goose	<i>Plectropterus gambensis</i>	Waterfowl	LC / LC	50
Red-eyed Dove	<i>Streptopelia semitorquata</i>		LC / LC	50
Three-banded Plover	<i>Charadrius tricollaris</i>		LC / LC	50
Cape Crow	<i>Corvus capensis</i>		LC / LC	50
Pied Starling	<i>Lamprotornis bicolor</i>	Gregarious	LC; SLS / LC	50
White-throated Swallow	<i>Hirundo albigularis</i>		LC / LC	50
Greater Striped Swallow	<i>Cecropis cucullata</i>		LC / LC	50
Cloud Cisticola	<i>Cisticola textrix</i>		LC; NE / LC	50
Wing-snapping Cisticola	<i>Cisticola ayresii</i>		LC / LC	50
Yellow-crowned Bishop	<i>Euplectes afer</i>	Gregarious	LC / LC	50
Speckled Mousebird	<i>Colius striatus</i>		LC / LC	45.5
Grey-winged Francolin	<i>Scleroptila afra</i>	Gamebird	LC; SLS / LC	37.5
Red-throated Wryneck	<i>Jynx ruficollis</i>		LC / LC	37.5
Diederik Cuckoo	<i>Chrysococcyx caprius</i>		LC / LC	37.5
Speckled Pigeon	<i>Columba guinea</i>		LC / LC	37.5
Laughing Dove	<i>Spilopelia senegalensis</i>		LC / LC	37.5
Spotted Thick-knee	<i>Burhinus capensis</i>	Gamebird	LC / LC	37.5
African Wattled Lapwing	<i>Vanellus senegallus</i>		LC / LC	37.5
Crowned Lapwing	<i>Vanellus coronatus</i>		LC / LC	37.5
Little Grebe	<i>Tachybaptus ruficollis</i>	Waterfowl	LC / LC	37.5
Bokmakierie	<i>Telophorus zeylonus</i>		LC / LC	37.5
Brown-throated Martin	<i>Riparia paludicola</i>		LC / LC	37.5
Zitting Cisticola	<i>Cisticola juncidis</i>		LC / LC	37.5
Pale-crowned Cisticola	<i>Cisticola cinnamomeus</i>		LC / LC	37.5
Cape Weaver	<i>Ploceus capensis</i>	Gregarious	LC; NE / LC	37.5
Red-billed Quelea	<i>Quelea quelea</i>	Gregarious	LC / LC	37.5
Pin-tailed Whydah	<i>Vidua macroura</i>	Gregarious	LC / LC	37.5
Giant Kingfisher	<i>Megaceryle maxima</i>		LC / LC	36.4

Common Name	Scientific Name	Priority Species	Threat Status (RSA / IUCN)	SABAP2 RR (%)
Pied Kingfisher	<i>Ceryle rudis</i>		LC / LC	36.4
White-breasted Cormorant	<i>Phalacrocorax lucidus</i>	Waterfowl	LC / LC	36.4
African Spoonbill	<i>Platalea alba</i>	Waterfowl	LC / LC	36.4
Secretarybird	<i>Sagittarius serpentarius</i>	SCC	VU / EN	30
Little Swift	<i>Apus affinis</i>	Gregarious	LC / LC	27.3
Black Sparrowhawk	<i>Accipiter melanoleucus</i>	Raptor	LC / LC	27.3
White-rumped Swift	<i>Apus caffer</i>	Gregarious	LC / LC	25
African Snipe	<i>Gallinago nigripennis</i>	Waterfowl	LC / LC	25
Common Greenshank	<i>Tringa nebularia</i>	Waterfowl	LC / LC	25
Common Buzzard	<i>Buteo buteo</i>	Raptor	LC / LC	25
Hamerkop	<i>Scopus umbretta</i>	Waterfowl	LC / LC	25
Glossy Ibis	<i>Plegadis falcinellus</i>	Waterfowl	LC / LC	25
Southern Bald Ibis	<i>Geronticus calvus</i>	SCC	VU; SLS / VU	25
White Stork	<i>Ciconia ciconia</i>	Gamebird	LC / LC	25
Pied Crow	<i>Corvus albus</i>		LC / LC	25
Cape Robin-Chat	<i>Cossypha caffra</i>		LC / LC	25
Cape Starling	<i>Lamprotornis nitens</i>		LC / LC	25
Common Myna	<i>Acridotheres tristis</i>		LC / LC	25
House Sparrow	<i>Passer domesticus</i>		LC / LC	25
Fan-tailed Widowbird	<i>Euplectes axillaris</i>	Gregarious	LC / LC	25
Black-throated Canary	<i>Crithagra atrogularis</i>	Gregarious	LC / LC	25
Rufous-naped Lark	<i>Mirafra africana</i>		LC / LC	20
South African Shelduck	<i>Tadorna cana</i>	Waterfowl	LC / LC	18.2
Red-billed Teal	<i>Anas erythrorhyncha</i>	Waterfowl	LC / LC	18.2
Western Barn Owl	<i>Tyto alba</i>	Raptor	LC / LC	18.2
Spotted Eagle-Owl	<i>Bubo africanus</i>	Raptor	LC / LC	18.2
Common Moorhen	<i>Gallinula chloropus</i>	Waterfowl	LC / LC	18.2
Common Sandpiper	<i>Actitis hypoleucos</i>		LC / LC	18.2
African Fish Eagle	<i>Haliaeetus vocifer</i>	Raptor	LC / LC	18.2
Jackal Buzzard	<i>Buteo rufofuscus</i>	Raptor	LC; NE / LC	18.2
Rock Martin	<i>Ptyonoprogne fuligula</i>	Gregarious	LC / LC	18.2
Dark-capped Bulbul	<i>Pycnonotus tricolor</i>		LC / LC	18.2
Cape White-eye	<i>Zosterops virens</i>		LC; NE / LC	18.2
Lesser Swamp Warbler	<i>Acrocephalus gracilirostris</i>		LC / LC	18.2
White-browed Sparrow-Weaver	<i>Plocepasser mahali</i>	Gregarious	LC / LC	18.2
Black-winged Lapwing	<i>Vanellus melanopterus</i>		LC / LC	16.7
Greater Kestrel	<i>Falco rupicoloides</i>	Raptor	LC / LC	16.7
Red-winged Starling	<i>Onychognathus morio</i>		LC / LC	16.7
Little Rush Warbler	<i>Bradypterus baboecala</i>		LC / LC	16.7
Common Quail	<i>Coturnix coturnix</i>	Gamebird	LC / LC	12.5
African Black Duck	<i>Anas sparsa</i>	Waterfowl	LC / LC	12.5
Cape Shoveler	<i>Spatula smithii</i>	Waterfowl	LC / LC	12.5
Black-collared Barbet	<i>Lybius torquatus</i>		LC / LC	12.5
Malachite Kingfisher	<i>Corythornis cristatus</i>		LC / LC	12.5
Marsh Owl	<i>Asio capensis</i>	Raptor	LC / LC	12.5

Common Name	Scientific Name	Priority Species	Threat Status (RSA / IUCN)	SABAP2 RR (%)
Blue Crane	<i>Grus paradisea</i>	SCC	NT / VU	12.5
Wood Sandpiper	<i>Tringa glareola</i>		LC / LC	12.5
African Darter	<i>Anhinga rufa</i>	Waterfowl	LC / LC	12.5
Grey Heron	<i>Ardea cinerea</i>	Waterfowl	LC / LC	12.5
Intermediate Egret	<i>Ardea intermedia</i>	Waterfowl	LC / LC	12.5
Black Stork	<i>Ciconia nigra</i>	SCC	VU / LC	12.5
Mountain Wheatear	<i>Myrmecocichla monticola</i>		LC / LC	12.5
Black-chested Prinia	<i>Prinia flavicans</i>		LC / LC	12.5
Willow Warbler	<i>Phylloscopus trochilus</i>		LC / LC	12.5
Eastern Clapper Lark	<i>Mirafra fasciolata</i>		LC / LC	12.5
Eastern Long-billed Lark	<i>Certhilauda semitorquata</i>		LC; SLS / LC	12.5
Spike-heeled Lark	<i>Chersomanes albofasciata</i>		LC / LC	12.5
Botha's Lark	<i>Spizocorys fringillaris</i>	SCC	EN; E / EN	12.5
Village Weaver	<i>Ploceus cucullatus</i>	Gregarious	LC / LC	12.5
Maccoa Duck	<i>Oxyura maccoa</i>	SCC	NT / EN	10
Southern Pochard	<i>Netta erythrophthalma</i>	Waterfowl	LC / LC	10
European Roller	<i>Coracias garrulus</i>	SCC	NT / LC	10
Grey Crowned Crane	<i>Balearica regulorum</i>	SCC	EN / EN	10
Ruff	<i>Calidris pugnax</i>		LC / LC	10
Martial Eagle	<i>Polemaetus bellicosus</i>	SCC	EN / EN	10
Rock Kestrel	<i>Falco rupicolus</i>	Raptor	LC / LC	10
Great Crested Grebe	<i>Podiceps cristatus</i>	Waterfowl	LC / LC	10
Goliath Heron	<i>Ardea goliath</i>	Waterfowl	LC / LC	10
Spotted Flycatcher	<i>Muscicapa striata</i>		LC / LC	10
Common Starling	<i>Sturnus vulgaris</i>		LC / LC	10
Orange-breasted Waxbill	<i>Amandava subflava</i>	Gregarious	LC / LC	10
Yellow Canary	<i>Crithagra flaviventris</i>	Gregarious	LC / LC	10
Golden-breasted Bunting	<i>Emberiza flaviventris</i>	Gregarious	LC / LC	10
Crested Barbet	<i>Trachyphonus vaillantii</i>		LC / LC	9.1
African Palm Swift	<i>Cypsiurus parvus</i>	Gregarious	LC / LC	9.1
Lanner Falcon	<i>Falco biarmicus</i>	SCC	VU / LC	9.1
Little Egret	<i>Egretta garzetta</i>	Waterfowl	LC / LC	9.1
Karoo Thrush	<i>Turdus smithi</i>		LC; NE / LC	9.1
Common House Martin	<i>Delichon urbicum</i>		LC / LC	9.1
Common Reed Warbler	<i>Acrocephalus baeticatus</i>		LC / LC	9.1
Great Reed Warbler	<i>Acrocephalus arundinaceus</i>		LC / LC	9.1
Fulvous Whistling Duck	<i>Dendrocygna bicolor</i>	Waterfowl	LC / LC	0
Black-winged Stilt	<i>Himantopus himantopus</i>		LC / LC	0
Pied Avocet	<i>Recurvirostra avosetta</i>		LC / LC	0
Whiskered Tern	<i>Chlidonias hybrida</i>		LC / LC	0
African Marsh Harrier	<i>Circus ranivorus</i>	SCC	EN / LC	0
Montagu's Harrier	<i>Circus pygargus</i>	Raptor	LC / LC	0
Lesser Kestrel	<i>Falco naumanni</i>	Raptor	LC / LC	0
Purple Heron	<i>Ardea purpurea</i>	Waterfowl	LC / LC	0
Greater Flamingo	<i>Phoenicopterus roseus</i>	SCC	NT / LC	0

Common Name	Scientific Name	Priority Species	Threat Status (RSA / IUCN)	SABAP2 RR (%)
Fiscal Flycatcher	<i>Melaenornis silens</i>		LC; NE / LC	0
Capped Wheatear	<i>Oenanthe pileata</i>		LC / LC	0
Red-collared Widowbird	<i>Euplectes ardens</i>	Gregarious	LC / LC	0

EN = Endangered; VU = Vulnerable; NT = Near Threatened; LC = Least Concern; E = Endemic to SA; NE = Near Endemic; SLS = Endemic to SA, Lesotho & Swaziland

## 4. FIELD RESULTS

### 4.1. SITE DESCRIPTION

The northern portion of the Sivutse B site is comprised of natural open grassland (~102 ha) making up approximately 33% of the site, while the southern section comprises mostly cultivated and fallow fields (~150 ha), making up approximately 49% of the site. Natural drainage lines with wetlands and a few small farm dams border the site on the northern and western side and in the southern section. A small stand of alien *Eucalyptus* trees occurs next to fallow fields in the north-eastern corner. A small rocky ridge occurs in the western corner of the site, where the remains of an old settlement is also evident. The grassland around the old structures is relatively disturbed and a few stands of alien *Eucalyptus* trees occur, mainly around the old farmstead. Secondary grassland on recovering old fields apparently associated with the old farmstead occur in the southwestern corner of the site. This small farm appears to have been active in the late 1950s according to historical aerial imagery (Figure 8).



Natural open grassland in the northern portion (left) and natural drainage line with wetlands and a small farm dam in the north-eastern corner (right) of the Sivutse B site



Cultivated fields in the southern portion (left) and fallow fields with a stand of alien trees on the eastern side (right) of the Sivutse B site



**Rocky ridge (left) and stand of alien trees around an old farmstead (right) in the western corner of the Sivutse B site**

The Sivutse A site is generally more disturbed by past activities (old Majuba Colliery underground mine), and is comprised primarily of disturbed grassland, with large stands of alien *Eucalyptus* trees in the highly disturbed areas. Parts of the site are completely modified by buildings (now derelict), roads, and other old infrastructure. A few drainage lines and wetlands occur around the site and a small dam occurs towards the centre (Figure 8).



**Disturbed grassland covering the majority of the Sivutse A site (top left); stands of alien trees and old infrastructure (top right); disturbed drainage line (bottom left); and small dam towards the centre of the site (bottom right)**




Figure 8: Habitat features of the Sivutse study area

#### 4.2. AVIFAUNAL HABITATS IN THE STUDY AREA AND SURROUNDS

The Sivutse B site and the surrounding areas are comprised of natural open grassland habitat and farmland, interspersed with natural watercourses with wetlands and small farm dams, that collectively create an ecosystem that supports many avifaunal species typical of the grassland biome. The most important habitat for avifauna occurring in the Sivutse study area is the natural open grassland vegetation, and the natural drainage lines occurring in the northern and eastern sections of the Sivutse B site. This natural grassland is connected to grasslands to the north and west. The extensive tracts of open grassland vegetation on this site and in the surrounding areas supports most of the terrestrial species found in the region, including priority species such as gamebirds, raptors, cranes, ibises, and smaller gregarious species. The many watercourses, farm dams, and wetlands provide important habitat for waterfowl and other wetland associated species such as ducks, geese, herons, and flamingos, while the wet areas provide surface water for drinking for all fauna. Another key habitat in the study area is the rocky ridge occurring in the western corner, which creates habitat heterogeneity within the landscape facilitating species diversity. Rocky areas provide a greater diversity of potential niches for plants and animals because of the microclimatic conditions they offer (Burnett *et al.*, 1998), therefore creating unique feeding opportunities in the landscape. Rocks also provide perching and display opportunities for many birds.



**Extensive natural open grassland in the northern section of the Sivutse B site**



**Small farm dam and open grassland typical of the surroundings just north of the Sivutse B site**




**Natural drainage line and wetlands in the northern section of the Sivutse B site**



**Rocky habitat in the western corner of the Sivutse B site**

The grassland vegetation on the Sivutse A site is more disturbed but does provide habitat for birds, although of a poorer quality. The small dam on the site also provides habitat for waterfowl and other wetland associated species.



**Small dam on the Sivutse A site**

#### **4.3. BIRD SPECIES OCCURRENCE IN THE STUDY AREA**

##### **4.3.1. BIRD OBSERVATIONS**

During the preliminary field survey, 55 species of birds were recorded in the study area and surroundings. Birds were identified either by direct observation (sighting and/or call) or by field signs such as tracks or feathers. These are listed in **Table 2** along with their national (Taylor *et al.*, 2015) and global (IUCN Red List of Threatened Species, 2023) conservation status. Bird species observed in the study area included mainly species typical of the grassland biome such as cisticolas, larks, longclaws, swallows, chats, and pipits. Birds recorded at the dams and wetlands included geese, ducks, herons, ibises, widowbirds, lapwings, and flamingos.

#### 4.3.2. BIRDS OF CONSERVATION CONCERN

Species of conservation concern (SCC) are those with a Red List status higher than Least Concern at a national level (Taylor *et al.*, 2015) and global level (IUCN, 2024) and/or species Protected at a national level (DFFE, 2023). SSC recorded during the preliminary site survey in the study area and surrounds are highlighted in red in **Table 2** below. These include Blue Crane *Grus paradisea* (NT / VU), Greater Flamingo *Phoenicopterus roseus* (NT / LC), and Southern Bald Ibis *Geronticus calvus* (VU / VU) recorded just to the north of the Sivutse B site and Southern Bald Ibis at the dam on the Sivutse A site. In addition, a nest of a Secretarybird *Sagittarius serpentarius* (VU / EN), in which the birds apparently successfully bred this past year, was recorded ~870 m to the north of the north-eastern corner of the Sivutse B site (see **Figure 8**).



Top to bottom: Blue Crane, Greater Flamingo, and Southern Bald Ibis recorded just north of the Sivutse B site



Southern Bald Ibis recorded at the dam on the Sivutse A site

#### 4.3.3. PRIORITY SPECIES

Preliminary assessment of species recorded in and around the study area show that there are some bird species that may be susceptible to the impacts of solar PV development occurring in the study area and surrounds. These include SCC as well as large-bodied, ground-welling species and waterfowl such as ducks, geese, flamingos, ibises, and cranes; raptors such as kites, buzzards, and Secretarybirds; and gregarious species such as sparrows, starlings, swallows, bishops, and widowbirds. These and other priority species recorded in the study area and surrounds are listed in **Table 2** along with their national and global conservation status.

**Table 2: Species recorded in the study area and surroundings listed in taxonomic order. SCC are highlighted in red**

Scientific Name	Common Name	National Status	Global Status	Priority Species
<i>Pternistis swainsonii</i>	Swainson's Spurfowl	LC	LC	Gamebird
<i>Coturnix coturnix</i>	Common Quail	LC	LC	Gamebird
<i>Numida meleagris</i>	Helmeted Guineafowl	LC	LC	Gamebird
<i>Alopochen aegyptiaca</i>	Egyptian Goose	LC	LC	Waterfowl
<i>Plectropterus gambensis</i>	Spur-winged Goose	LC	LC	Waterfowl
<i>Anas undulata</i>	Yellow-billed Duck	LC	LC	Waterfowl
<i>Lybius torquatus</i>	Black-collared Barbet	LC	LC	
<i>Ceryle rudis</i>	Pied Kingfisher	LC	LC	
<i>Columba livia</i>	Rock Dove	LC	LC	
<i>Columba guinea</i>	Speckled Pigeon	LC	LC	
<i>Streptopelia capicola</i>	Ring-necked Dove	LC	LC	
<i>Grus paradisea</i>	Blue Crane	NT	VU	SCC
<i>Gallinula chloropus</i>	Common Moorhen	LC	LC	Waterfowl
<i>Burhinus capensis</i>	Spotted Thick-knee	LC	LC	Gamebird
<i>Himantopus himantopus</i>	Black-winged Stilt	LC	LC	Waterfowl
<i>Vanellus armatus</i>	Blacksmith Lapwing	LC	LC	Waterfowl
<i>Vanellus senegallus</i>	African Wattled Lapwing	LC	LC	Gamebird
<i>Elanus caeruleus</i>	Black-winged Kite	LC	LC	Raptor
<i>Buteo buteo</i>	Common Buzzard	LC	LC	Raptor
<i>Sagittarius serpentarius</i>	Secretarybird	VU	EN	SCC
<i>Falco rupicoloides</i>	Greater Kestrel	LC	LC	Raptor
<i>Microcarbo africanus</i>	Reed Cormorant	LC	LC	Waterfowl
<i>Ardea melanocephala</i>	Black-headed Heron	LC	LC	Gamebird
<i>Bubulcus ibis</i>	Western Cattle Egret	LC	LC	Gamebird
<i>Phoenicopterus roseus</i>	Greater Flamingo	NT	LC	SCC

Scientific Name	Common Name	National Status	Global Status	Priority Species
<i>Plegadis falcinellus</i>	Glossy Ibis	LC	LC	Waterfowl
<i>Bostrychia hagedash</i>	Hadada Ibis	LC	LC	Gamebird
<i>Geronticus calvus</i>	Southern Bald Ibis	VU	VU	SCC
<i>Threskiornis aethiopicus</i>	African Sacred Ibis	LC	LC	Gamebird
<i>Lanius collaris</i>	Southern Fiscal	LC	LC	
<i>Corvus capensis</i>	Cape Crow	LC	LC	
<i>Telophorus zeylonus</i>	Bokmakierie	LC	LC	
<i>Cossypha caffra</i>	Cape Robin-Chat	LC	LC	
<i>Saxicola torquatus</i>	African Stonechat	LC	LC	
<i>Myrmecocichla formicivora</i>	Ant-eating Chat	LC	LC	
<i>Lamprotornis bicolor</i>	Pied Starling	LC	LC	Gregarious
<i>Hirundo rustica</i>	Barn Swallow	LC	LC	Gregarious
<i>Cecropis cucullata</i>	Greater Striped Swallow	LC	LC	Gregarious
<i>Petrochelidon spilodera</i>	South African Cliff Swallow	LC	LC	Gregarious
<i>Cisticola tinniens</i>	Levaillant's Cisticola	LC	LC	
<i>Cisticola juncidis</i>	Zitting Cisticola	LC	LC	
<i>Cisticola textrix</i>	Cloud Cisticola	LC	LC	
<i>Cisticola cinnamomeus</i>	Pale-crowned Cisticola	LC	LC	
<i>Cisticola ayresii</i>	Wing-snapping Cisticola	LC	LC	
<i>Acrocephalus gracilirostris</i>	Lesser Swamp Warbler	LC	LC	
<i>Calandrella cinerea</i>	Red-capped Lark	LC	LC	
<i>Passer melanurus</i>	Cape Sparrow	LC	LC	Gregarious
<i>Motacilla capensis</i>	Cape Wagtail	LC	LC	
<i>Macronyx capensis</i>	Cape Longclaw	LC	LC	
<i>Anthus cinnamomeus</i>	African Pipit	LC	LC	
<i>Ploceus velatus</i>	Southern Masked Weaver	LC	LC	Gregarious
<i>Euplectes orix</i>	Southern Red Bishop	LC	LC	Gregarious
<i>Euplectes progne</i>	Long-tailed Widowbird	LC	LC	Gregarious
<i>Estrilda astrild</i>	Common Waxbill	LC	LC	Gregarious
<i>Vidua macroura</i>	Pin-tailed Whydah	LC	LC	Gregarious

EN = Endangered; VU = Vulnerable; NT = Near Threatened; LC = Least Concern

#### 4.4. KEY HABITATS AND PRELIMINARY SITE SENSITIVITY

The extensive open grassland vegetation in the surroundings, with natural drainage lines, wetlands, and dams, provide the main habitats that support the avifaunal species found in the region, including priority species such as cranes, Secretarybirds, ibises, flamingos, raptors, and gregarious species. Approximately 102 ha of natural open grassland vegetation occurs in the northern section of the Sivutse B site and is relatively undisturbed. According to historical aerial imagery (from ~1957 to today) this area has not been modified by ploughing in the last ~67 years and has likely only been used for grazing. The natural grassland vegetation as well as drainage lines, wetlands, and dams in the Sivutse study area have been rated as highly sensitivity due to the natural extent and intactness (high functional integrity), as well as the support of SCC (conservation value). The rocky ridge in the western portion is also rated as highly sensitive as this area provides habitat heterogeneity facilitating higher biodiversity and provides specialised habitat in the landscape. The disturbed grassland on the Sivutse A site provides relatively poor grassland habitat for birds and has a medium-low sensitivity rating. Fallow fields and stands of alien trees have a low sensitivity rating and areas that have been completely modified such as cultivated fields, buildings and roads have a very-low sensitivity rating (Figure 9).

The relatively undisturbed areas should be avoided by the proposed development. The natural grassland should be avoided as far as possible, while the drainage lines and wetland areas must be avoided by the buffer specified by the wetland and/or aquatic specialists. From a species perspective, a preliminary buffer of 1 km has been applied to the Secretarybird nest following preliminary consultation with the relevant authorities (EWT Birds of Prey Programme and BirdLife SA). The nest occurs ~870 m from the north-eastern corner of the Sivutse B site, and a small portion of the site (~3.5 ha) falls within this buffer (see **Figure 9**). The effectiveness of the buffer will need to be investigated further during the follow-up seasonal monitoring and in the Avifaunal Impact Assessment to follow. In addition, according to BirdLife SA, large, contiguous habitat for Secretarybirds that has high functional integrity (i.e. habitat remnants of 100 ha or more) should be avoided by development (BirdLife SA, 2022). Refer to **Figure 9** for the preliminary assessment of sensitivity for the site features represented in **Figure 8** with recommendations following the Species Environmental Assessment Guideline (SANBI, 2020) and summarised in **Table 3**.

**Table 3: Summary of sensitivity categories**

Site Feature	Description and Recommendation	Sensitivity Rating
Natural open grassland	Natural grassland provides the main habitat for the avifauna, including SCC, found in the area, especially Secretarybird. Large, contiguous patches of grassland ( $\geq 100$ ha) should be avoided. Avoidance mitigation wherever possible. Minimisation mitigation – changes to project infrastructure design to limit the amount of habitat impacted; limited development activities of low impact acceptable. <b>Offset mitigation may be required</b> for high impact activities.	High
Watercourses / drainage lines / wetlands / farm dams	Water courses / drainage lines and natural wetlands or small farm dams provide important aquatic habitat for waterfowl and other priority species. These features must be avoided by the development by the buffer specified by the wetland / aquatic specialist/s.	High
Rocky ridge	Rocky ridges and outcrops are areas of high biodiversity and provide specialist habitat in the landscape. Avoidance mitigation wherever possible. Minimisation mitigation – changes to project infrastructure design to limit the amount of habitat impacted; limited development activities of low impact acceptable. <b>Offset mitigation may be required</b> for high impact activities.	High
Secretarybird Nest	A preliminary buffer of 1 km has been applied to the nest following preliminary consultation with the relevant authorities ( <b>Figure 9</b> ). This buffer may be revised following further investigation.	High
Disturbed wetlands and channels	Wetlands and drainage lines on the Sivutse A site that are either highly disturbed or modified (e.g. channel). Minimisation and restoration mitigation applies – development activities of medium impact followed by appropriate restoration activities.	Medium
Disturbed Grassland	Disturbed grassland on the Sivutse A site. Provides secondary habitat for avifauna. Minimisation and restoration mitigation applies – development activities of medium impact followed by appropriate restoration activities.	Medium-low
Modified and disturbed areas – stands of alien trees and fallow fields	Areas that have been modified by alien trees and past cultivation. Little to no natural vegetation occurs in these areas. These areas do not need to be avoided by the development, unless a nest of a priority species is recorded in the follow-up surveys during the pre-construction monitoring.	Low
Modified areas – cultivated fields, buildings, and roads	Areas that have been modified by cultivation, buildings, and roads. No natural vegetation occurs in these areas. These areas do not need to be avoided by the development.	Very Low




Figure 9: Preliminary avifaunal habitat sensitivity of the Sivutse study area

## 5. SITE SENSITIVITY VERIFICATION

The Renewstable<sup>®</sup> Sivutse B site is comprised of approximately 102 ha of natural open grassland vegetation that forms a large continuous expanse with the surrounding areas, and represents the most important habitat for birds in the landscape, including many SCC, and is rated as **highly sensitive**. Other key habitats that are considered highly sensitive include drainage lines, wetlands, farm dams, and the rocky ridge in the western corner. As such, the nature of the landscape supports SCC such as Blue Crane, Greater Flamingo, Secretarybird, and Southern Bald Ibis. For the Animal Species theme for the Sivutse B site, the sensitivity rating identified by the screening tool of **High** for Aves corresponds with the large expanse of natural grassland habitat and freshwater ecosystems in the study area and is appropriate.

The grassland vegetation on the Sivutse A site is disturbed and represents poorer / secondary grassland habitat for birds. The majority of the site is rated as **medium sensitivity**. For the Animal Species theme for the Sivutse A site, the sensitivity rating identified by the screening tool of **High** for Aves is less appropriate and can drop to **Medium**.

During the preliminary field surveys, a number of bird SSC were recorded in the study area and surroundings, including the Sivutse A site. This includes two species listed by the Screening Tool Report, Southern Bald Ibis *Geronticus calvus* (VU) and Secretarybird *Sagittarius serpentarius* (VU). Therefore, once the development footprints have been finalised, a full assessment of potential impacts that the proposed development may impose on avifauna, should be conducted for both sites following the Species Environmental Assessment Guidelines (SANBI, 2020) in accordance with the *Protocol for the Specialist Assessment and Minimum Report Content Requirements for Environmental Impacts on Animal Species* (GN 1150 of 30 October 2020) as amended 28 July 2023.

## 6. PRELIMINARY IMPACTS

The overall environmental impacts of solar energy developments are poorly understood globally. Unlike wind energy developments, there is presently no clear pattern in the types of birds negatively affected by solar plants, and collision casualties recorded to date include a wide variety of avian guilds (Jenkins *et al.*, 2017). Widely accepted impacts of solar PV include permanent habitat destruction, fragmentation, and the associated bird displacement (particularly for range restricted species), as well as collision with reflective panels as birds mistake large panel arrays for wetlands, otherwise known as the “lake effect” (Lovich and Ennen, 2011; Smit, 2012; DeVault *et al.*, 2014; Visser, 2016; Kosciuch *et al.*, 2020; Chock *et al.*, 2021). Other general impacts documented to date include noise and disturbance caused by construction activities, attraction of novel species through the creation of artificial nest sites and shade, and chemical pollution from panel cleaning (Lovich and Ennen, 2011; DeVault *et al.*, 2014; Chock *et al.*, 2021). The impacts of additional infrastructure associated with solar energy developments, such as roads, power lines, and substations, must also be considered. These include, habitat destruction, fragmentation, threat of collision, and electrocution (Jenkins *et al.*, 2017).

Possible impacts on avifauna during the construction and operational phases and their sources associated with the proposed development are provided in **Table 4**. The installation of the proposed Renewstable<sup>®</sup> Sivutse project and ancillary infrastructure will require clearance of a relatively large amount of natural grassland vegetation during the construction phase. The majority of the site will be comprised of solar PV arrays during the operational phase. The main impact relating to avifauna will therefore be loss of some natural habitat, the displacement of species including gregarious species. Other possible direct impacts include possible collisions of priority species moving through the area with panels and power lines during the operational phase. Possible

indirect impacts include spread of invasive alien vegetation due to disturbance to the soil, and potential contamination of the soil and downstream watercourses should chemicals be used to clean the panels.

**Table 4: Possible impacts arising from the proposed development**

Possible Impact	Source of Impact	Area and Species to be Affected	Development Phase	Nature of Impact
Loss of vegetation and avifaunal habitat	Clearing vegetation for installation of infrastructure including solar panels, roads, and buildings	Natural open grassland (Sivutse B site); Terrestrial grassland species; large-bodied, ground-dwelling gamebirds including SCC, raptors	Construction	Direct
Collision of avifauna with reflective surfaces of solar panels leading to injury or death	Solar panels perceived to be water body by avifauna	Solar PV development site; Gamebirds, waterfowl; raptors; SCC such as cranes and Secretarybirds	Operation	Direct
Collision and/or electrocution of avifauna with associated power lines	Power lines	Power line route; Gamebirds, waterfowl; raptors; SCC such as cranes and Secretarybirds	Operation	Direct
Contamination of the environment by hazardous materials	Cleaning of solar panels during operation	Solar PV development site; All species	Construction and Operation	Indirect
Spread of invasive alien plant species	Disturbance to soil and clearing of vegetation	Study area and surroundings	Construction	Indirect
Disturbance and displacement of resident bird species	Clearing of site and construction activities; Operational and maintenance activities; attraction of novel species	Site and immediate surroundings; Small terrestrial species; Ground-dwelling gamebirds; SCC such as cranes and Secretarybirds	Construction and Operation	Indirect
Increased human disturbance; Gradual environmental degradation	<ul style="list-style-type: none"> <li>Disturbance to the study area, adding to existing pressures in the landscape</li> <li>Adding to cumulative pressures in the landscape caused by other approved or proposed renewable energy projects</li> </ul>	Study area and surrounding natural areas	Operation	Cumulative

## 7. PRE-CONSTRUCTION MONITORING REQUIREMENTS

The Renewable<sup>®</sup> Sivutse study area supports many priority species (bird species that may be susceptible to the impacts of solar PV development), including SCC such as Blue Crane, Greater Flamingo, Secretarybird, and Southern Bald Ibis. The BirdLife SA guidance notes on minimising the impacts of infrastructure development on Secretarybirds *Sagittarius serpentarius* (BirdLife SA, 2022) states that large, contiguous patches of grassland (>= 100 ha) should be avoided. In addition, a recently used Secretarybird nest was recorded ~870 m to the north of the Sivutse B site, to which a preliminary development buffer of 1 km has been applied.

It is therefore important to conduct pre-construction monitoring according to the Best Practice Guidelines for Birds and Solar Energy (Jenkins *et al.*, 2017) for assessing and monitoring the impact of solar power generating facilities on birds in southern Africa. This will take the avifaunal assessment to **Stage 2 – Data Collection**, which includes structured and repeated data collection on which to base the impact assessment report and provide a baseline against which post-construction monitoring can be compared.

The duration and scope of data collection is guided by the size of the proposed development as well as the results of the preliminary assessment, which verifies the sensitivity of avifauna potentially affected by the proposed development (see **Table 5**). For the Sivutse B site, assessment **Regime 2** would be the minimum requirement (see **large** solar facilities >150 ha / >50 MW) with the current layout, and sampling over **two** seasons would need to be conducted, with one survey falling within the peak (summer) season. If the Secretarybirds are found to be breeding during this time, then it is recommended that the monitoring regime be extended to **three** seasons. For the Sivutse A site, assessment **Regime 2** would also be applicable (see **medium** solar facilities 30-150 ha) with the current layout with sampling conducted over **two** seasons, with one survey falling within the peak (summer) season.

**Table 5: Recommended avifaunal assessment regimes (Jenkins *et al.*, 2017)**

Type	Size	Avifaunal Sensitivity*		
		Low	Medium	High
All solar technologies except Concentrated Solar Power (CSP)	Small (<30 ha / <10 MW)	<b>Regime 1</b> One site visit of 1-5 days	<b>Regime 1</b> One site visit of 1-5 days	<b>Regime 2</b> 2-3 seasonal visits of 3-5 days over 6 months Pre- & post-con monitoring mortality searches
	Medium (30-150 ha / 10-50 MW)	<b>Regime 1</b> One site visit of 1-5 days	<b>Regime 2</b> 2-3 seasonal visits of 3-5 days over 6 months Pre- & post-con monitoring mortality searches	<b>Regime 2</b> 2-3 seasonal visits of 3-5 days over 6 months Pre- & post-con monitoring mortality searches
	Large (>150 ha / >50 MW)	<b>Regime 2</b> 2-3 seasonal visits of 3-5 days over 6 months Pre- & post-con monitoring mortality searches	<b>Regime 2</b> 2-3 seasonal visits of 3-5 days over 6 months Pre- & post-con monitoring mortality searches	<b>Regime 3</b> 4-5 seasonal visits of 4-8 days over 12 months Pre- & post-con monitoring mortality searches
CSP	All	<b>Regime 3</b> 4-5 seasonal visits of 4-8 days over 12 months Pre- & post-con monitoring mortality searches		

\* The avifaunal sensitivity is based on the number of priority species present, or potentially present, the regional, national, or global importance of the affected area for these species (both individually and collectively), and the perceived susceptibility of these species (both individually and collectively) to the anticipated impacts of development

Based on the key habitats observed in the study area and surrounds, the following sampling would need to be incorporated into the data collection for each season:

- Abundance estimates for small terrestrial birds through point count or walked linear transect surveys.
- Counts for large terrestrial birds and raptors in the study area and surrounds, through driven road transects and vantage point monitoring.

- Observations of flight behaviour of priority species flying over or near the proposed development area and associated risk of collision.
- Counts of bird numbers at focal wetlands such as the farm dams and local movements between waterbodies.
- Searches for nest sites of large terrestrial species and any habitats likely to support nest sites of key raptors and other priority species should be surveyed and checked on each survey to confirm occupancy. Any evidence of breeding activity and/or its outcomes must be recorded.
- Surveys of existing nearby power lines for signs of bird collisions and electrocutions.
- Details of any incidental sightings of priority species.

## 8. RECOMMENDATIONS

The following recommendations are intended to guide the positioning of the proposed infrastructure and layout:

- All natural drainage lines, wetlands, and dams, as well as rocky ledges, must be avoided, including the buffer recommended by the aquatic and/or wetland specialist/s.
- A preliminary buffer of 1 km has been applied to the Secretarybird nest. It is important that the infrastructure remain out of the buffer to avoid disturbance to potentially breeding birds, especially during the construction phase.
- It will be important to conduct pre-construction monitoring, especially on the Sivutse B site, due to the potential impacts on the Secretarybird and other priority species.
- For both sites, it is advisable that assessment **Regime 2** covering **two seasons** be followed for the pre-construction monitoring.
- If the Secretarybirds are found to be breeding during this time, then it is recommended that the monitoring regime for the Sivutse B site be extended to **three** seasons.

## 9. REFERENCES

- BirdLife SA (2022): *Minimising the Impacts of Infrastructure Development on Secretarybirds Sagittarius serpentarius*, Guidance Note, 06 October 2022, BirdLife SA.
- Chock, R.Y., Clucas, B., Peterson, E.K., *et al.* (2021): Evaluating potential effects of solar power facilities on wildlife from an animal behavior perspective, *Conservation Science and Practice*, 2021;3:e319. <https://doi.org/10.1111/csp2.319>
- Coordinated Waterbird Counts (CWAC): <https://cwac.birdmap.africa/sites.php?sitecode=31421812>
- DEA (2018): National Protected Areas Expansion Strategy (NPAES) for South Africa, Pretoria: Department of Environmental Affairs.
- DeVault, T.L., Seamans, T.W., Schmidt, J.A., Belant, J.L., Blackwell, B.F., Mooers, N., Tyson, L.A. and Van Pelt, L. (2014): Bird use of solar photovoltaic installations at US airports: implications for aviation safety, *Landscape and Urban Planning* 122: 122-128.
- DFFE (2024a): South African Protected Area Database (SAPAD), Pretoria: Department of Forestry, Fisheries and the Environment, [https://egis.environment.gov.za/protected\\_areas\\_register](https://egis.environment.gov.za/protected_areas_register)
- DFFE (2024b): South African Conservation Area Database (SACAD), Pretoria: Department of Forestry, Fisheries and the Environment, [https://egis.environment.gov.za/protected\\_areas\\_register](https://egis.environment.gov.za/protected_areas_register)
- Gill, F., Donsker, D., and Rasmussen, P. (Eds). (2023): *IOC World Bird List (v13.2)*. doi: 10.14344/IOC.ML.13.2.

- Global Biodiversity Information Facility (GBIF): Free and open access to biodiversity data: <https://www.gbif.org/>
- Harrison, J.A., Allan D.G., Underhill, L.G., Herremans, M., Tree, A.J., Parker, V. and Brown, C.J. (1997): *The Atlas of Southern African Birds*, Johannesburg: BirdLife South Africa.
- Hockey, P.A.R., Dean, W.R.J. and Ryan P.G. (2005): *Roberts Birds of Southern Africa, 7th Edition*, Cape Town: John Voelcker Bird Book Fund.
- iNaturalist (California Academy of Sciences and the National Geographic Society) Online Database: <https://www.inaturalist.org/>
- IUCN (2024): IUCN Red List of Threatened Species, Version 2024-1: <http://www.iucnredlist.org>
- Jenkins, A.R., Ralston, S. and Smit-Robinson, H.A. (2017): *Birds and Solar Energy Best Practice Guidelines: Best Practice Guidelines for assessing and monitoring the impact of solar power generating facilities on birds in southern Africa*, Johannesburg: BirdLife South Africa.
- Kosciuch, K., Riser-Espinoza, D., Geringer, M., Erickson, W. (2020): A summary of bird mortality at photovoltaic utility scale solar facilities in the Southwestern U.S., PLoS ONE 15(4): e0232034.
- Lovich, J.E. and Ennen, J.R. (2011): Wildlife conservation and solar energy development in the desert southwest, United States, *BioScience* 61: 982-992.
- Marnewick, M.D., Retief, E.F., Theron, N.T., Wright, D.R., and Anderson, T.A. (2015): *Important Bird and Biodiversity Areas of South Africa*, Johannesburg: BirdLife South Africa.
- Mucina, L. and Rutherford, M.C. (2006): The vegetation of South Africa, Lesotho and Swaziland, *Strelitzia* 19, Pretoria: South African National Biodiversity Institute.
- SANBI (2020): Species Environmental Assessment Guideline. Guidelines for the implementation of the Terrestrial Fauna and Terrestrial Flora Species Protocols for environmental impact assessments in South Africa, Pretoria: South African National Biodiversity Institute, Version 3.1. 2022.
- South African Bird Atlas Project 2 (SABAP2) (2024): <http://sabap2.adu.org.za/index.php/>
- Southern African Birding cc. (2011): Roberts VII Multimedia, Birds of Southern Africa, Computer Software.
- Taylor, M.R., Peacock, F. and Wanless, R.M. (eds.) (2015): *Eskom Red Data Book of Birds of South Africa Lesotho and Swaziland*, Johannesburg: BirdLife South Africa.
- Taylor, M.R., Peacock, F. and Wanless, R.M. (eds.) (2015): *Eskom Red Data Book of Birds of South Africa Lesotho and Swaziland*, Johannesburg: BirdLife South Africa.
- Visser, E. (2016): *The impact of South Africa's largest photovoltaic solar energy facility on birds in the Northern Cape, South Africa*, Masters Dissertation, University of Cape Town.

## 10. APPENDICES

### APPENDIX A: ABRIDGED CV OF THE SPECIALIST

Name and Surname	:	Robyn Phillips
Date of Birth	:	28 08 1975
Company Name	:	Cossypha Ecological
Field of Expertise	:	Terrestrial Ecologist and Avifaunal Specialist
SACNASP Registration	:	<i>Pr.Sci.Nat.</i> 400401/12 (Zoological and Ecological Sciences)
Highest Qualification	:	MSc (Zoology) <i>cum laude</i>
Years of Experience	:	23
Contact Number	:	084 695 1648
Email	:	robyn@cosypha.co.za

The first half of my professional career was spent working in ecological research at the University of KwaZulu-Natal. Since starting in consulting in 2011, I have been involved in many projects requiring biodiversity surveys and ecological assessments as part of the legislated requirements for the Environmental Impact Assessment (EIA) process. These studies include field assessment of habitat, species occurrence (especially those of conservation concern), assessment of ecological importance and sensitivity of floral and faunal communities and habitat, as well as assessment of impacts. Tasks also include making recommendations and prescribing mitigation measures after applying the mitigation hierarchy, aimed at minimising impacts.

Following is a selection of similar projects undertaken:

- Avifaunal Impact Assessments for the proposed Bateleur Solar PV Cluster Development between Mopane and Musina, Limpopo Province (ABO Wind) – 2023 to present.
- Avifaunal Impact Assessments for the proposed Kwena Solar PV Cluster Development near Groblersdal, Limpopo Province (ABO Wind) – 2023 to present.
- Avifaunal Impact Assessments for the proposed ZCC N3 Solar PV Developments along the N3 from Ashburton to Heidelberg (EnviroAfrica) – 2023 to present.
- Avifaunal Impact Assessments for the proposed Nyala Solar PV Developments near Northam, Limpopo Province (PRAXOS 373) – 2022 to present.
- Avifaunal Impact Assessments for the proposed Ndau Solar PV Developments near Polokwane, Limpopo Province (PRAXOS 373) – 2022 to present.
- Avifaunal Impact Assessment for the proposed Harvard 1 & 2 Solar PV Plants and Grid Connection, Bloemfontein, Free State (EnviroAfrica) – 2021 to 2023.
- Terrestrial Biodiversity and Faunal Assessment for the proposed Springhaas Solar Cluster Development and Grid Connection near Dealesville, Free State (GIBB Environmental) – 2021 to 2023.
- Avifaunal Impact Assessment, Terrestrial Fauna Compliance Statement, and Terrestrial Biodiversity Impact Assessment for the proposed Oceana 10 MW Solar PV Facility near St Helena Bay, Western Cape (SRK) – 2021 to 2022.
- Terrestrial Biodiversity (including fauna and flora) and Avifaunal Impact Assessment for the Waterkloof Solar IPP Programme, North West (GIBB Environmental) – 2020 to 2021.
- Avifaunal Assessment for the Proposed Development of a Battery Energy Storage System (BESS) and Associated Infrastructure at the Cuprum Substation located at Copperton, near the town of Prieska, Northern Cape Province (AECOM) – 2021.
- Terrestrial Biodiversity Assessment (including flora, fauna, and avifauna) for the Askham Solar Energy Facility, Northern Cape (Komani San) – 2018 to 2019.
- Faunal and Avifaunal Assessments for various solar farms in the Northern Cape (SEF) – 2011 to 2012.